

Applications of the Mellin-Perron Formula in Number Theory: Addendum Number II: asymptotics of the coefficients of a Dirichlet series

Marko R. Riedel

October 30, 2022

1 Introduction

This addendum to my MSc. thesis treats the following problem: given the closed form of a Dirichlet series, how can we recover the asymptotics of the coefficients of the series?

There are two parts to this document: first, we apply generalized Mellin summation to obtain an integral formula for the coefficients and second, we use this to compute the asymptotics of the sum-of-divisors function.

References to my thesis will be provided throughout.

2 Generalized Mellin summation

We will be using generalized Mellin summation (harmonic sums) as described in section 2.10, page 87 of the thesis.

The function $f(x)$ is given by

$$f(x) = \delta(x - n)$$

where $\delta(x)$ is the Dirac delta function and n is a positive integer and its Mellin transform is

$$f^*(s) = \int_0^{+\infty} f(x)x^{s-1}dx = n^{s-1}.$$

We take $F(x) = \sum_k \lambda_k f(\mu_k x)$ with $\mu_k = k$ and let $\Lambda(s) = \sum_k \frac{\lambda(k)}{k^s}$. The Mellin transform of $F(x)$ is

$$F^*(s) = \Lambda(s)n^{s-1}.$$

Applying Mellin inversion now yields

$$F(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Lambda(s)n^{s-1}x^{-s}ds.$$

Setting $x = 1$, we find

$$F(1) = \sum_k \lambda_k \delta(k - n) = \lambda_n$$

and hence

$$\lambda_n = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Lambda(s)n^{s-1}ds.$$

3 Asymptotics of the sum-of-divisors function

Let

$$\sigma_k(n) = \sum_{d|n} d^k \quad \text{and} \quad \Lambda_k(s) = \sum_n \frac{\sigma_k(n)}{n^s}.$$

Now $\sigma_0(n) = d(n)$. This immediately implies that

$$\Lambda_0(n) = \zeta^2(s).$$

In order to compute the asymptotics of the sum-of-divisors function σ_k we need a closed form of its Dirichlet series.

3.1 Dirichlet series of σ_k , $k > 0$

$$\zeta(s)\zeta(s-k) = \sum_{p=1}^{\infty} \frac{1}{p^s} \sum_{q=1}^{\infty} \frac{q^k}{q^s} = \sum_{n=1}^{\infty} \sum_{pq=n} \frac{1}{p^s} \frac{q^k}{q^s} = \sum_{n=1}^{\infty} \sum_{q|n} \frac{q^k}{n^s} = \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q|n} q^k = \Lambda_k(s).$$

3.2 Asymptotics of the sum-of-divisors function

We have

$$\operatorname{Res} [\zeta^2(s)n^{s-1}; s = 1] = \ln n + 2\gamma$$

and hence

$$d(n) \sim \ln n + 2\gamma.$$

Furthermore for $k > 0$, we have

$$\operatorname{Res} [\zeta(s)\zeta(s-k)n^{s-1}; s = k + 1] = \zeta(k + 1)n^k$$

and

$$\operatorname{Res} [\zeta(s)\zeta(s-k)n^{s-1}; s = 1] = \zeta(1 - k)$$

and hence

$$\sigma_k(n) \sim \zeta(k + 1)n^k + \zeta(1 - k).$$

The first few values are

$$\begin{aligned}\sigma_1(n) &\sim \frac{\pi^2}{6}n - \frac{1}{2} \\ \sigma_2(n) &\sim \zeta(3)n^2 - \frac{1}{12} \\ \sigma_3(n) &\sim \frac{\pi^4}{90}n^3 \\ \sigma_4(n) &\sim \zeta(5)n^4 + \frac{1}{120} \\ \sigma_5(n) &\sim \frac{\pi^6}{945}n^5\end{aligned}$$

4 External links

- Marko Riedel <http://pnp.mathematik.uni-stuttgart.de/iadm/Riedel/index.html> *Applications of the Mellin-Perron Formula in Number Theory.*