Dweak squares in Heller triangulated categories

Matthias Künzer

November 5, 2007

Abstract

We mention briefly an elementary compatibility of distinguished weak (dweak) squares with the shift functor, which holds in a closed Heller triangulated category.

In a Verdier triangulated category, one can still formulate this compatibility assertion, but I do not know whether it can be proven there.

To obtain a counterexample to this compatibility assertion in a Verdier triangulated category, however, one would need a Verdier triangulated category that is not a closed Heller triangulated category. I do not know an example of such a category.

Let $(\mathcal{C}, \mathsf{T}, \vartheta)$ be a closed Heller triangulated category; cf. [**2,** Def. 1.5.i], [**3,** Def. A.6]. Recall that closedness just means that each morphism can be completed to a 2-triangle. Heller triangulated categories in which idempotents split, are closed; cf. [**2,** Lem. 3.1]. The stable category of a Frobenius category is closed.

We use the conventions and notations of [**2**].

A closed Heller triangulated category is in particular Verdier triangulated, taking the set ([1]) of 2-triangles as set of distinguished triangles; cf. [**5,** Def. 1-1]; this is proven just as [**2,** Prop. 3.6].

Suppose given $n \geqslant 2$. Each chain of morphisms $X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-1}} X_n$ in $\mathcal{C}$ can be prolonged to an n-triangle, and this prolongation is unique up to isomorphism; cf. [**2,** Def. 1.5.ii.2, Lem. 3.1, Lem. 3.4.6].

A *distinguished weak square*, for short *dweak square* ([2]), is a weak square whose diagonal sequence fits into a 2-triangle. Dweak squares are indicated by the symbol Ξ. Note that to define dweak squares, a Verdier triangulated category suffices.

Since a completion of a morphism to a 2-triangle is unique up to isomorphism, so is completion of an angle to a dweak square  ; and dually.

Using a 4-triangle, one sees that dweak squares compose. There is an elegant method, due to Neeman, to show this fact already in Verdier triangulated categories; cf. [**4,** Lem. 2.1].

[1] In a sufficiently big universe.

[2] Also known as a *homotopy cartesian square.*

Completing iteratively to dweak squares, we may therefore form



and the resulting chain $X'_1 \xrightarrow{f'_1} \cdots \xrightarrow{f'_{n-1}} X'_n$ is unique up to isomorphism.

Remark 1. *The following compatibility of dweak squares and shift holds in our Heller triangulated category* $(\mathcal{C}, \mathsf{T}, \vartheta)$.

$$(X'_1 \xrightarrow{f'_1} \cdots \xrightarrow{f'_{n-1}} X'_n) \simeq (X_1^{+1} \xrightarrow{f_1^{+1}} \cdots \xrightarrow{f_{n-1}^{+1}} X_n^{+1}) .$$

Proof. Complete the chain $X_1 \xrightarrow{f_1} \cdots \xrightarrow{f_{n-1}} X_n$ to an n-triangle. By [**2,** Lem. 3.4.1, Lem. 3.4.2], an n-triangle consists of dweak squares. The result follows from the uniqueness up to isomorphism stated just before Remark 1. ▫

In a Verdier triangulated category, in which the axiom [**1,** 1.1.13] holds, the assertion of Remark 1 holds if

$$n = 2$$

and, by the octahedral axiom in the form of loc. cit., if

$$n = 3 .$$

But we may ask for this assertion for

$$n \geqslant 4$$

as well. In fact, both sides of the isomorphism in question are still welldefined up to isomorphism resp. welldefined.

If $n = 4$, the valid assertion in the cases $n = 2$ and $n = 3$, together with [**2,** Lem. 3.4.1], yields isomorphisms

$$\begin{array}{ccc} X'_1 & \xrightarrow{f'_1} & X'_2 \\ {\scriptstyle u_1^{1,2}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_2^{1,2}} \\ X_1^{+1} & \xrightarrow{f_1^{+1}} & X_2^{+1} \end{array} \qquad \begin{array}{ccc} X'_2 & \xrightarrow{f'_2} & X'_3 \\ {\scriptstyle u_2^{2,3}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_3^{2,3}} \\ X_2^{+1} & \xrightarrow{f_2^{+1}} & X_3^{+1} \end{array} \qquad \begin{array}{ccc} X'_1 & \xrightarrow{f'_1 f'_2} & X'_3 \\ {\scriptstyle u_1^{1,3}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_3^{1,3}} \\ X_1^{+1} & \xrightarrow{f_1^{+1} f_2^{+1}} & X_3^{+1} \end{array}$$

$$\begin{array}{ccc} X'_1 & \xrightarrow{f'_1 f'_2 f'_3} & X'_4 \\ {\scriptstyle u_1^{1,4}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_4^{1,4}} \\ X_1^{+1} & \xrightarrow{f_1^{+1} f_2^{+1} f_3^{+1}} & X_4^{+1} \end{array} \qquad \begin{array}{ccc} X'_2 & \xrightarrow{f'_2 f'_3} & X'_4 \\ {\scriptstyle u_2^{2,4}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_4^{2,4}} \\ X_2^{+1} & \xrightarrow{f_2^{+1} f_3^{+1}} & X_4^{+1} \end{array} \qquad \begin{array}{ccc} X'_3 & \xrightarrow{f'_3} & X'_4 \\ {\scriptstyle u_3^{3,4}}\downarrow\wr & & \wr\downarrow{\scriptstyle u_4^{3,4}} \\ X_3^{+1} & \xrightarrow{f_3^{+1}} & X_4^{+1} \end{array}$$



But in a Verdier triangulated category with **[1,** 1.1.13] added, I do not know how to prove that, say, $u_3^{1,2,3}$ may be chosen equal to $u_3^{3,4}$.

Neither do I know a Verdier triangulated category in which the assertion of Remark 1 fails, say, for $n = 4$. Worse still, I do not know an example of a Verdier triangulated category that is not closedly Heller triangulated.

References

[1] Bernstein, J.; Beilinson, A.A.; Deligne, P., *Faisceaux pervers*, Astérisque 100, 1982.

[2] Künzer, M., *Heller triangulated categories,* Homol., Homot. Appl. 9 (2), p. 233–320, 2007.

[3] Künzer, M., *Nonisomorphic Verdier octahedra on the same base,* preprint, arxiv, 0708.0151, 2007.

[4] Neeman, A., *K-theory for triangulated categories I (A),* Asian J. Math. 1 (2), p. 330–417, 1997.

[5] Verdier, J.L., *Catégories Derivées*, published in SGA 4 1/2, SLN 569, p. 262–311, 1977 (written 1963).

MSC2000: 18E30.